


Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

Reserve

A99.24
L72Ae

UNITED STATES
DEPARTMENT OF AGRICULTURE
LIBRARY



Reserve

BOOK NUMBER


942314

A99.24
L72Ae

3. NOTES ON TROPICAL DENDROLOGY //

BY

ELBERT L. LITTLE, JR.
Forester (Dendrology)
United States Forest Service
Washington, D.C., U. S. A.



CONTENTS

	<u>Page</u>
Introduction	1
The nomenclature of trees	3
The classification of trees	6
The identification of trees	9
The collection of botanical specimens	12
The herbarium	16
Botanical terms	17
The distribution of trees	22
Ecology	23

Prepared for the Tropical Forestry Training Course held March to May 1955 at the Tropical Forest Research Center of the Forest Service, United States Department of Agriculture, at Río Piedras, Puerto Rico, with the cooperation of the Foreign Operations Administration.

These notes are a slightly revised translation of the author's *Notas sobre Dendrología Tropical*, mimeographed in 1953 while the author was visiting professor of dendrology at the forestry school of Universidad de los Andes, Mérida Venezuela. It is planned to expand these notes into a reference on tropical dendrology to include additional information on the important families of tropical forest trees.

United States Forest Service
San Río Piedras, Puerto Rico
1955

942814
INTRODUCTION

What is dendrology? The term dendrology (dendrologia in Spanish) is a scientific and artificial word. It is derived from two Greek words meaning trees and discourse or study, or the study of trees. The limits of the science of dendrology are controversial. A review of the history of usage of the term has been made by William A. Dayton (What is dendrology? Journal of Forestry 43:710-722. 1945).

Perhaps the first use of the word dendrology was in the year 1668 as the title of a book or encyclopedia on trees by Ulisse Aldrovandi, Italian physician and naturalist. Originally dendrology included all aspects of trees, and in that time there was no science of forestry. Now and especially in Europe dendrology includes also the shrubs, but in the United States of America frequently it is restricted to trees.

In actual use, dendrology is limited to the botany of trees or precisely to the taxonomy of trees. It can be considered as a division of forestry or botany which treats of the taxonomy of trees. In some universities of the United States the subject matter of dendrology is taught by a professor of forestry and in others by a professor of botany who is a specialist in taxonomy or systematic botany.

Dendrology, then, is a division of forestry or botany which treats of the taxonomy of trees and other woody plants, including nomenclature, classification, identification, and distribution.

The subject matter in tropical countries should be called tropical dendrology, or the taxonomy of tropical trees, in order to distinguish it from a course in dendrology as taught in a university of the United States or Europe. Those countries and continents of the temperate zone have trees very different from those of tropical regions. A forester who has studied dendrology only in a forestry school in the United States knows very little about tropical trees, for example.

Why study dendrology?--There are five principal objectives in the study of tropical dendrology:

1. Nomenclature of trees. To learn how trees are named, including scientific names, common names, and the code of botanical nomenclature.

2. Classification of trees. To learn how trees are classified or arranged into families, genera, and other groups according to their arrangements. To learn the names and characteristics of the common and important botanical families of trees.

3. Identification of trees. To be able to place an unknown tree in its family. To learn how to find the name of unknown trees or to identify trees by means of keys, manuals, and floras. To know the reference books for identification of the trees of your country. To learn how to collect botanical specimens. To learn how to maintain and use a herbarium.

4. Distribution of trees. To learn how trees are distributed into climatic zones and forest types. To know the geographic distribution of important forest trees.

5. Important forest trees. To know the important forest trees of your country, including scientific name, common names, family, distribution and abundance, and uses.

Reference books on dendrology.--There are very few textbooks on dendrology. In the United States of America there are a few, such as Textbook of Dendrology by William M. Harlow and Ellwood S. Harrar (Ed. 3, 555 pages, illustrated. 1950). Also various manuals and floras of regions and States are used. However, the books do not include tropical trees and are not useful in the tropical countries.

As there are no texts of tropical dendrology for the tropical American countries, these notes may serve as an introduction to the subject. References and texts of plant taxonomy have much material that relates to dendrology. Many of these are published in English and other languages, rather than Spanish. Various books, pamphlets, and articles on the trees, woods, and plants of a country or locality are useful to foresters in that country.

What is a tree?--Everybody knows what a tree is, but it is not easy to prepare a precise definition. It is possible to classify seed plants, or flowering plants, into four artificial groups on the basis of size and habit of the stems: trees, shrubs, herbs, and vines. The ancient Greek Theophrastus (372-287 B. C.), disciple of Aristotle and called the father of botany, distinguished trees, shrubs, and herbs. These artificial groups are not related to the natural classification of botany into the botanical families.

The definitions in Forest Terminology (by the Society of American Foresters, 1944; translated into Spanish by M. A. González Vale as terminología Forestal, 1950) are modified slightly here.

Tree (árbol).--A woody plant with a well defined erect perennial trunk and a more or less definitely formed crown and which generally attains a height of at least 12 to 15 feet (or 4 or 5 meters) and a trunk diameter of 3 or 4 inches (7 to 10 centimeters) at breast height (4 1/2 feet or 1.3 meters).

Shrub (arbusto).--A perennial woody plant smaller than a tree, usually with several perennial stems branched from the base.

Herb (hierba).--A plant with a herbaceous or soft stem, annual or perennial, but not woody. (A herb can be annual or perennial or in cold climates with a stem that dies to the ground each year.)

Vine (bejuco).--A woody or herbaceous plant with the stems not erect but depending on other plants or objects for support.

Practical applications of dendrology.--Dendrology is like a tool or an instrument for becoming acquainted with and studying the trees. The names serve as a guide for referring to the trees. Before making a survey of the forest resources of a tropical country, it is necessary to know the names of the species. Always foresters need to know the names of the important trees with which they work.

THE NOMENCLATURE OF TREES

Nomenclature is a division of taxonomy which treats of the names of plants, including the correct names, synonyms, and rules of nomenclature.

Trees, like other plants, have two kinds of names, common names and scientific names. Both are important and necessary and both have their advantages and disadvantages.

Common names are in the language of the country. Their origin is an interesting subject. Some Spanish names of trees in tropical America are words used before by the American Indians. Others are the same as those of different trees in Spain. Perhaps the Spanish colonists gave these names to unknown trees of the New World which resemble trees of the Old World. Some common names are of other kinds, such as descriptive, from the use, from the environment, or from the region or place where the species is found. In the United States other English common names are translations of the scientific name or in honor of the discoverer or another person.

Advantages of common names.- 1. They are in the language known by the people. 2. They are used by the country persons, woodsmen, people in general, and in commerce.

Disadvantages of common names.--1. They change in different places, countries, and languages.

2. The same common names can be used for different species in different places, countries, etc.

3. Many species do not have their own distinct common names. There are still unknown species without names.

4. Many common names are not exact. Some species have indefinite common names which correspond only to a genus or to a botanical family.

5. There is no definite authority nor code of rules governing common names and for making them uniform.

As they are in modern languages, common names are useful only in one language and change from one country to another. A useful tree species of extensive distribution can have five to ten or more names in various localities and in commerce. For example, a tree of the West Indies can have an English name in Jamaica, a Spanish name in Cuba, and a French name in Haiti. And perhaps in the other islands, such as Puerto Rico and the Lesser Antilles it has other names. If it extends to South America or Central America, the names can change more. Also, there is confusion because the same common names can be used for different species in various parts of its natural range.

Then, for exactness and clarity and to avoid confusion, the botanists and also the foresters are obliged to use the scientific names of trees.

Advantages of scientific names.--1. They are uniform in a universal system in use throughout the world.

2. They are in the Latin language, which is the language of no country and being a dead language does not change through the years.

3. There is the International Code of Botanical Nomenclature of rules for scientific names and for naming new species.

4. They show the classification and relationships of the species.

Disadvantages of scientific names.--1. They are strange and long.
2. They are not used by the people.

The Latin language which was used by the scholars of past centuries was continued by the biologists for the scientific names of plants and animals. A few centuries ago the botanists were studying medicinal plants or herbs and used to write books with the descriptions and text in Latin. The Latin description of a phrase served as the name.

Carolus Linnaeus (1707-1778), the distinguished Swedish naturalist, established the binomial system of nomenclature in the year 1753. In that year he published in Latin his book *Species Plantarum* (Species of Plants), which is the beginning of modern botanical nomenclature.

The binomial system of nomenclature, or system of two names, means that the name of each species of plants consists of two Latin words, the genus and the specific epithet. (The same system is used for animals.)

For example, the scientific name of the species of mahogany of Central America and South America is *Swietenia macrophylla*. To these two words the systematic botanists add the name of the author, the botanist who first gave this name to the species and published a botanical description of it. Thus, *Swietenia macrophylla* King. Botanical works should include the author's name. However, generally it is unnecessary to write nor remember the author, and foresters seldom need to mention the author.

Scientific names are subject to definite rules. These rules are adopted and revised by the systematic botanists in international botanical congresses. The last congresses were at Stockholm, Sweden, in 1950, and at Paris, France, in 1954. The most recent edition of the rules, published in 1952, is called International Code of Botanical Nomenclature, by J. Lanjouw and others (228 pages). Minor changes or amendments were made in 1954 and will be incorporated in a revised edition.

Under the Code the scientific names are in Latin or if from other languages or artificial origin, they have Latin endings. The generic name is a noun and begins with a capital letter. The specific epithet commences with a small letter and can be: (1) an adjective which agrees with the generic name in gender (masculine, feminine, or neuter); (2) a noun in the Latin genitive case such as the name of a person; or (3) the name of another genus or another plant in apposition.

The two words are underlined in manuscripts or on the typewriter and are in italics in publications. In origin and derivation the scientific names are descriptive or of other kinds like the common names.

There are three very important rules in the Code. The rule of types states that a scientific name is based upon a specimen called the type. The identity is fixed with this specimen which is kept in a large herbarium. According to the rule of priority, the correct name of a group is the oldest which is in accord with the Code. For example, in the past many species have been given more than one scientific name by different botanists working independently. Thus, there is only one valid name, the oldest, and the others are called synonyms. The rule of homonyms treats of homonyms or identical names. The same name cannot be used for two different groups, and if a name was used earlier for one group, it never can be employed for another.

From these rules it may be seen that scientific names are not perfect. Among the botanists there is no complete agreement in the names, their application, and their limits. Some species still have two scientific names in use in different books. However, the scientific names are much more distinct and clearer than the common names.

Pronunciation of scientific names.--Being in Latin, the scientific names are pronounced like Latin words. The Latin language is pronounced almost like Spanish, and the vowels are similar in both languages. Then, it is simple and easy in Spanish speaking countries to treat the scientific names as if they were written in Spanish. However, there are some names derived from foreign words, such as modern languages. These names are pronounced as in the original language, often with accent different from Spanish. The British and other Europeans pronounce scientific names correctly according to Latin, but in the United States these words generally are spoken as if English.

Abbreviation of names of authors.--The names of some authors are written in abbreviated form after the scientific names. Generally these are botanists who have named many species or who have long names.

Complete names of these persons can be found in glossaries of some botanical references. Usually the abbreviation stops just before the second vowel. An exception is that of Carolus Linnaeus, only L. For example, Rhizophora mangle L., mangrove or mangle. Also, there is seen H. B. K., for Humboldt, Bonpland, and Kunth, as in Byrsonima crassifolia H. B. K., chaparro.

Double citation of names of authors.--Some scientific names of plants are followed by names of two authors, the first in parenthesis. For example, Delonix regia (Bojer) Raf., flamboyant-tree or flamboyán. This means that the first author gave the name of the specific epithet but in another genus or as a variety. Afterwards, the second author changed the name and put the specific epithet in this arrangement. In this case the earlier name, which also is now in use, is Poinciana regia Bojer. Some botanists regard Delonix as a genus distinct from Poinciana and others do not.

THE CLASSIFICATION OF TREES

Classification is a division of taxonomy which treats of the botanical arrangement of plants into groups, such as families and genera in accord with the relationships.

This is the problem. There are approximately 350,000 known species of living plants. It is not possible to study and know them all one by one. How can they be arranged into groups for study, for compilation of data on the characteristics, and for organization of all this information?

There are two kinds, artificial classifications and natural classifications. An artificial classification is a simple and convenient arrangement but not according to the relationships. It is like compartments or pigeon-holes in a box or cabinet, one compartment for each species.

The ancient Greek Theophrastus proposed the artificial classification previously mentioned. This arrangement of plants on the basis of habit of stems as trees, shrubs, and herbs is useful and convenient. Foresters study mainly the trees, which form an artificial group.

Another artificial classification was the sexual system of Carolus Linnaeus, published in 1732. All plants were placed in 24 classes based upon the stamens, their number, union, and length. The classes were divided into orders, based upon the number of the styles in each flower. This system served to identify specimens and was very useful in its time.

A natural classification attempts to group together similar plants according to their relationships. After Linnaeus, other botanists proposed natural systems of classification of plants. In these works the species were arranged in natural groups such as the families. The French botanist Antoine de Jussieu made one of the first natural systems in the year 1789.

The modern classification of plants and animals is based upon the principle or theory of organic evolution. In 1859 the British naturalist Charles Darwin published his famous work. The Origin of Species. The principle of organic evolution means simply that the higher forms of plants and animals have developed from the simple or lower forms over millions and millions of years. Specialized plants have originated from primitive species. In other words, plant life has changed slowly during long periods of time.

Natural classification is based upon relationships through descent. Evolution can be compared with a tree. In theory plant life began as a seed. Through millions of years it grew into a tree with many branches representing the plant kingdom. The buds corresponds to the species that exist now, and the branches to the extinct or fossil species. Then, all the twigs on one branch belong to the same family and are related. But as the branches do not exist now, the relationships are not well known and are subject to differences of opinion among botanists.

There are many evidences and proofs to support the principle of organic evolution. Morphology, or the study and comparison of the form and parts of plants, is important. For example, those species with similar form or structure are thought to be related. Other evidences can be found in other subdivisions of biology such as anatomy, embryology, genetics, cytology, paleontology, and geographical distribution.

The methods of organic evolution are not so well understood. Among the theories is the theory of mutation (or of sudden changes in the hereditary variations) and the theory of natural selection (or survival of the fittest) by Darwin.

Probably the natural system of classification of plants most generally adopted by botanists at present is that of Engler and Prantl, two German botanists, in their important work of 20 volumes entitled *Die Natürlichen Pflanzenfamilien* (1887 and following years; The Natural Plant Families), which covers the entire plant kingdom. The most recent evidences indicate that perhaps this system could be improved. Nevertheless, it is the most detailed and most convenient and is used in many large herbaria of the world.

Another important natural system also in use is that of Bentham and Hooker, two British Botanists, in their Latin work of 3 volumes, *Genera Plantarum* (1862-63; The Genera of Plants).

The categories of the plant kingdom.--In the natural classification, the species of trees and other plants are arranged into groups of small and large rank in a hierarchy. These groups of the plant kingdom are classed categories. The categories are in Latin and also in modern languages. They are listed below in Latin, English and Spanish, with examples.

Latin	English	Spanish
Regnum Vegetabile	Plant Kingdom	Reino Vegetal
Divisio	Division	División
Classis	Class	Clase
Ordo	Order	Orden
Familia	Family	Familia
Genus	Genus	Género
Species	Species	Especie
(Varietas)	(Variety)	(Variedad)
Example in Latin	Example in English	Example in Spanish
Spermatophyta (Divisio)	Spermatophytes (Division)	Espermatofitas
(Subdivisio Angiospermae)	(Subdivision Angiosperms)	(Angiospermas)
Dicotyledoneae	Dicotyledons	Dicotiledóneas
Geraniales	Geraniales	Geraniales
Meliaceae	Mahogany family	Meliáceas
Swietenia	Mahogany	Cuoba
macrophylla	Central American	de Honduras

At the end but not a category is the individual (individuum in Latin and individuo in Spanish). Also subgroups for other categories can be added in large groups as needed, such as subdivision in the example above, subfamily, subgenus, etc.

The plant kingdom now consists of approximately 350,000 known species of living plants grouped into 19,000 genera. The division of Spermatophytes (phanerogams or seed plants) now contains 2 subdivisions, 7 classes, 45 or more orders, more than 300 families, more than 10,000 genera, and more than 250,000 species.

Then, the most important unit in the botanical classification is the species. Each individual, tree or other plant, belongs to a species and only to one particular species. It is difficult to define a species and also the other categories. It can be said that the species is composed of individual plants (or animals) which are similar in appearance and which can reproduce or breed among themselves and produce other individuals resembling the parents.

A genus is a group of related species. A family also is composed of a group of related genera. An order consists of a group of related families, etc.

The variety is a division or minor variation of a species or a group of individuals which differ slightly from the others. The majority of species have no varieties or are not divided into varieties. Varieties are named particularly in cultivated species.

Scientific names of families and higher categories are plural, while names of genera, species and varieties are singular.

The name of an order ends in -ales and is derived from its type family. For example, Geraniales is from the family Geraniaceae, which is derived from the genus Geranium.

The termination of names of the botanical families is -aceae. However, the Code permits the use of eight exceptions with endings in -ae. For example Palmae, Leguminosae, and Guttiferae. The name of a family is derived from its type genus or from a synonym. For example, Meliaceae comes from Melia.

The subject matter of tropical dendrology includes under classification the study of families of the important forest trees with distinguishing characteristics and examples.

THE IDENTIFICATION OF TREES

The identification of a tree consists in determining the correct scientific name, generally by means of manuals, floras, keys, etc., or in determining that the plant or specimen is the same as a previously known plant with a scientific name. In these references there is a special botanical terminology for describing the differences in morphology or in the parts of the trees. For this reason, in the laboratory we study the terminology of the leaf, flower, fruit, etc.

Methods of identification of trees.--The question is: How to learn the name of a tree? There are several methods. Then, in each case we should use the easiest, simplest, and most rapid method which also arrives at the correct name.

1. The question.--The simplest method of learning the name of a tree is to ask someone who knows the name. This method can be used anywhere. Whenever there is the opportunity, we should go to the forests with other foresters or botanists who know the species well. This method is very useful, particularly in a new region where many trees are strange. In the university and in the herbarium, as in the field, the question aids identification.

This method is especially important for learning common names, because many common names are not found in the books. The rural people who know well the trees of their locality have learned the names from other persons and not from botanical books. When in doubt, one should ask two persons to check if both give the same name. Also, when the common name is known, it is easy frequently to obtain the scientific name of the genus or species in references on plants or woods.

However, there are limitations and disadvantages in the method of questions. (1) Other persons, including specialists can be mistaken in the names and in the identifications. (2) In some localities there are no persons who know all the trees, especially the scientific names. (3) Many times foresters have to work alone and where there is no help in the identifications. Therefore, forests need to know how to identify trees and botanical specimens as well.

2. Books, manuals, floras, catalogs, keys, monographs.--Wherever there is a good illustrated manual of the region, as in various parts of the United States, one can look through the illustrations. This method, useful though not scientific, can lose time and cannot be employed where there are many tree species, an illustrated manual could not illustrate many species of minor importance.

These books generally are written by botanists for botanists and with the technical terminology of systematic botany. Thus, foresters in the study of dendrology should learn to read and understand these botanical books, which have numerous, perhaps too many, technical terms. There is a need for more popular illustrated manuals containing a minimum of technical terms and written for foresters and the public.

Therefore, we shall study in the laboratory the botanical terminology of the leaf, the flower, the fruit, and other parts of the trees, such as the stem and the bark.

A flora of a region generally contains botanical descriptions and keys. However, some tropical countries lack descriptive floras.

A catalog has a list of the species of a region, often with other notes. The Catálogo de la Flora Venezuela also has keys to genera.

A monograph is a study of a genus or family in a country or larger region. For example, Rubiaceae of Venezuela by Standley and Podocarpus in the New World by Buchholz and Gray.

A key, like a key to a door, is a simple device for opening the way to the name, or an artificial device for finding rapidly the scientific name of a plant. This is much easier than reading many descriptions. The old botanical references of one or two centuries ago had no keys. In order to identify an unknown plant with a botanical book without a key, it is necessary to read the descriptions until reaching one which agrees with the plant. Thus, it is necessary to read half the book on an average in the identification of a specimen.

The key is dichotomous or with forks or branches two by two. It divides the plants of a book into groups of two or by halves until it arrives at the name which corresponds to the specimen. In a key there are pairs of contrasting short phrases, generally of a single line each. It is necessary to determine which of the two phrases agrees with the specimen. If the phrase contains two or more parts, all characters should agree with the specimen. Below the correct phrase is found another pair of contradictory phrases. The selection of a correct phrase is repeated until one arrives at the name. If there is a description, one should read it in order to check whether it fits the specimen. If it does not agree, probably there is an error and use of the key should be repeated by hunting another fork which leads to the correct identification.

There are keys to families and to genera within a family and to species within a genus. But unfortunately, in some tropical regions there are few keys to species. When there are two or more keys for use, it is simplest to use the shortest or that of the smallest region or with the smallest number of parts.

The two kinds of keys are the indented key and the parallel key. The indented key begins with the contrasting phrases of the pair at the left of the page usually not together but separated by other pairs following a little more to the right. Generally there are numbers or letters to facilitate the comparison but not in some short keys. Below each line the next pair is indented.

The parallel key always has the two contrasting phrases of the pair together, one directly below the other and the position of the pairs which follow is indicated by numbers.

We should know how to use both kinds of keys, because both are found in reference books. Each has its advantages and disadvantages. The long keys of many pages usually are parallel and do not lose space as in an indented key with many short lines. The indented key generally is preferred, because it is easier to follow, and if an error is made, the correct branch can be found more rapidly.

3. The herbarium.--A herbarium is a collection of plant specimens, dried, pressed, mounted in cardboards, identified, and arranged according to a botanical classification. The specimens are placed in large cases or cabinets of steel or wood. Special ones are preserved by other methods such as large fruits in boxes and succulent plants or small plants in liquid.

The herbarium serves for the identification of botanical specimens and also for reference and teaching. Type specimens are the base of botanical nomenclature, and all the specimens together show the geographic distribution. The herbarium is the basis of many investigations and publications of taxonomy.

How is the herbarium used in the identification of trees? There are two principal methods.

1. To verify the identification made by a key or a description in a botanical reference. It is much easier to compare a specimen with a named specimen than with a description.

2. For comparison with specimens of various species. When there are no botanical references adequate for identification, the herbarium is very useful. For example, in a country which lacks a descriptive flora or keys to species, it is necessary to take the botanical specimens to a herbarium that has collections from the country or the region in order to find identified specimens of the same species.

Whenever the herbarium is used in identification, first the botanical references should be consulted. It saves time to identify the specimen as completely as possibly with keys and floras before. Time is lost in using the herbarium before eliminating many similar species with the books. The genus of most plants can be determined with the books of other regions.

If the specimen is poor or is not complete with flowers and fruits, it is difficult to follow a key. Then the herbarium can be examined by trial and error in search for identical specimens, but this method is slow.

In the identification of a dried specimen in the herbarium, frequently it is necessary to dissect the flower in order to study the parts. The dried flower can be soaked in hot or boiling water before examination. Then it will become soft and can be examined without breaking and nearly as well as when fresh.

4. Shipment of specimens to a large herbarium.--Instead of making his own determinations, the forester can send botanical specimens to a large herbarium for identification.

In order to facilitate the determinations and also to prepare specimens that will be preserved permanently in a large herbarium, foresters should know how to collect and prepare good specimens.

The principal advantage of sending specimens to a large herbarium is that the identifications by the specialists will be correct and in accord with the most recent studies.

The disadvantages are that it is necessary to collect and prepare botanical specimens and that frequently there is a delay of some months or more than a year until the determinations are received. (Generally small collections are named faster than large ones.)

Before shipping the specimens for identification, one should always write and mention the number of specimens and the region or State where they were collected. Some herbaria do not have specialists in plants of certain parts of the world and are not interested in specimens from those countries. The service of identifications is free, but in place of payment the herbarium keeps the specimens.

When there is a specialist working on a particular group, such as a family, and writing monographs on its genera, this botanist is the best authority on that group and can make the best identifications.

Many countries have national herbaria which vary greatly in size. The largest herbaria in the world are principally in Europe and North America. Extensive collections from tropical America are deposited in three herbaria in the United States: Chicago Natural History Museum, Chicago, Illinois; New York Botanical Garden, Bronx Park, New York City, New York; and United States National Museum, Washington 25, D.C. Specialists from these and other herbaria in the United States have made extensive botanical explorations and collected many thousands of specimens in tropical America.

THE COLLECTION OF BOTANICAL SPECIMENS

The necessity of collecting botanical specimens.--In the identification of an unknown tree a specimen is worth more than many words or notes. It is difficult to identify a tree solely from the notes written in the forest.

From time to time foresters need to collect some botanical specimens of the trees with which they work. If they do not know the trees in the forest, then they should preserve specimens for identification later or for shipment to a large herbarium or to a specialist for determination.

For example, on making an inventory of the forest resources of a region, numbers or common names can be used for the unknown trees. Then, specimens should be collected for later identification.

Systematic botanists have various methods of collecting specimens. However, foresters generally collect only a few specimens and can employ the simple methods. In an emergency a twig can be broken from a tree and pressed in a notebook, in the pocket, or in a book. But it is worth the trouble to collect good specimens and in the end the identifications will result better.

Instructions for making collections. Collect sufficient material of good specimens with flowers or fruits.--This is the first rule. Frequently botanists are interested in collecting incomplete specimens of rare plants, but foresters are not. A rare tree generally is unimportant in forestry.

It is very difficult to identify correctly and completely a sterile specimen of a tree from only the leaves and twigs. Trees in various families have very similar leaves. Nevertheless, often unknown trees are found with neither flowers or fruits, and it is necessary to collect sterile specimens. Possibly there can be found a tree flowering outside the regular season. If not, it is very useful to collect a wood sample from the same tree. Perhaps specialists in wood anatomy can identify the genus or family of the wood, and then botanists can continue the identification.

Collect at least two sets.--Two specimens should be collected from the same tree, one to keep and the other to send to the herbarium or specialist for identification. Also a few additional flowers for dissection will be helpful. Whenever convenient, it would be useful to collect more sets. However, more time is spent and more equipment is needed in preparing the additional sets. In order to avoid a mixture of two species, all sets should be from the same tree.

Write useful notes in the forest for the label.--A specimen without notes is not worth much either to the forester or to the herbarium. In the herbarium the notes on the label probably are worth more than the specimen. These notes include:

1. Name of the collector.
2. Number of the specimen. Each collector should use a series of numbers for convenience in his notes, in identification, and in the herbarium when later reference is made to the specimens.
3. Date.
4. Locality. The country, State, municipality, exact locality or distance and direction from a city or from a point on the map.
5. Elevation above sea level.
6. Forest type. Associated species. Soil. Or if planted rather than wild.
7. Common name.
8. Size and habit. Height and diameter of the tree, or if the plant is a shrub, vine, or herb.

9. Uses.

10. Other notes. Abundance. Color of the flower. Fruit. Bark and latex. Wood. Roots.

Naturally all these notes cannot be obtained for all specimens.

Collecting equipment.--It is difficult to collect botanical specimens of trees because the branches usually are high. In a search there may be located a small tree of the same species or a tree with low limbs on the edge of the forest. Sometimes the tree can be climbed or felled. Or leaves, flowers, and fruits fallen on the ground beneath can be found.

In obtaining specimens, tools such as knives, machetes, and axes are indispensable. Also pruning shears and a pruning hook or pole may be used.

A notebook with pencil or pen, a hand lens, and a ruler or measuring tape are useful. So are field glasses and a camera.

It is important to carry the specimens from the forest to the office or base without losing parts, without damage, and without drying. Probably the best method is to carry a light press with old newspapers directly to the forest. Or the specimens can be placed in a canvas bag or wrapped in newspapers. Also, there are vasculums or large cans that are used with moist newspapers inside. For large fruits and seeds paper sacks are useful.

Pressing equipment.--Botanical specimens in the herbarium are dried and pressed, in order to preserve better and save space. The problem in drying the specimens is to remove the moisture with pressure and rapidly. If it is not pressed, the specimen does not stay flat but wrinkles and folds. Then it is fragile and breaks and becomes damaged easily. If it is not dried rapidly and completely, mold damages it. Also, when dried rapidly the specimens retain their natural color better and do not lose leaves or other parts.

In an emergency specimens can be pressed solely with some newspapers and a weight such as books, boards, bricks, or stones.

The press.--Generally a wood press size 12 by 17 or 18 inches with two straps of leather or canvas is used.

The old newspapers or newsprint which are folded for the specimens should be the same size as the press. Also 50 to 100 or more sheets of blotting paper or driers such as of felt are needed.

The sample is arranged within the newspaper. It should not be larger than the cardboard mounting sheets, which measure 16 1/2 by 11 1/2 inches. A large specimen can be folded in the form of V, N, or M. Since the mounting sheet and pressed specimen have only two dimensions instead of three, some leaves and twigs should be removed, leaving the base of the petioles to indicate the leaf arrangement. Generally one should not arrange a leaf directly on top of another. If too many leaves are left, the specimen does not dry well and the leaves covered beneath cannot be seen.

Outside on the corner of the newspaper sheet can be placed the number of the specimen which corresponds to the number of the collector in his notebook. Between each two folded papers containing specimens is inserted a drier. When all the specimens are arranged in papers, the press is tightened and fastened.

The press should be put in a warm dry place. Daily or twice daily the driers are changed until the specimens within the papers become completely dry and break when they are bent. The first time that the driers are changed the specimens should be examined and rearranged as needed, those that have become folded being straightened. The moist driers are dried in the sun, near a stove, or placed one by one against the walls and on the floor of a room. If driers are lacking, newspapers can be utilized. The time for drying varies from a few days up to a week, depending upon the size of the specimens, thickness of the leaves, etc. This method is rather slow and consumes much time and does not prepare good specimens of trees with thick leaves. However, it serves well enough for foresters who do not collect many specimens and do not wish to carry much equipment.

Heat.--Botanists who collect botanical specimens in quantities in tropical regions always use heat to dry the specimens rapidly. In the laboratory electric ovens with ventilators or fans can be used or an apparatus of electric heaters or electric light bulbs under the press.

In the field a portable stove of gasoline or kerosene or kerosene lanterns can be employed.

Corrugated driers.--With heat there are needed corrugated sheets or ventilators of cardboard or metal in order that the hot air can pass within the press and the water leave the specimens. The corrugated sheets have the same size of the press. Those of cardboard can be flat on one or both sides. The metal sheets are made of aluminum or iron and last more. They fit together in less space and dry the specimens in half the time required with cardboard. After 12 hours or more in an ordinary press with driers, the corrugated sheets are inserted in place of every second drier, and the press is mounted above the stove. A cloth is placed around the stove and fastened tightly around the bottom of the press but opened near the bottom or one side to permit entrance of air. With metal sheets the majority of the specimens will dry within 12 to 24 hours. At about the middle of this time the straps should be tightened and the press turned over. As there is danger of fire, the stove never should be left unattended.

After being dried the specimens should be arranged in order by number and kept in cardboard boxes or packages and in a dry place. If for much time, insecticides should be applied.

THE HERBARIUM

From time to time foresters collect botanical specimens for large herbaria and also use the herbaria in identification. Perhaps with the duplicate specimens they may make small herbaria of the regions where they work. Therefore, they should know something of the arrangement and care of the herbarium. How to identify specimens in the herbarium was discussed earlier.

The mounting of specimens.--After being identified and before being placed in the steel or wood cabinets, the specimens are mounted on mounting sheets of white cardboard. These cardboards which measure 16 1/2 by 11 1/2 inches, should be of good, durable quality. In some herbaria the specimens are mounted with glue or paste. In others narrow pieces of gummed cloth tape are used. With both methods the thick parts, such as stout twigs and large fruits, can be sewed with needle and thread. Loose parts, such as additional flowers and fruits are placed in pockets or envelopes glued to the mounting sheet.

The label.--The label is glued or pasted on the lower right corner of the mounting sheet. Generally labels are printed in part. The notes by the collector should be added with a typewriter. In addition there are added the scientific name with author, the name of the person who made the identification, and preferably also the year of the identification.

Genus folders.--The mounted specimens of a genus or of a species if there are many, are placed in a folded cardboard sheet 12 by 17 1/4 inches after folding. The generic name is placed on the lower left corner outside.

Arrangement of the herbarium.--Large herbaria generally follow a natural system of classification in their arrangement. The most popular is that of Dalla Torre and Harms, based upon that of Engler and Prantl. and with numbers for the families and genera. Each genus of seed plants has a number, and the specimens are arranged and may be found in this numerical order.

In a small herbarium of a region or in a personal herbarium it is simpler to follow the order of the botanical flora of the country. Generally the order of the families in botanical references is that of Engler and Prantl. Within a family it is simplest to arrange the genera and species alphabetically.

Protection against insects and mold.--It is always necessary to protect the herbarium against insects and in humid climates also from mold. No method is perfect. Perhaps the simplest is to place repellents, such as paradichlorobenzene or naphthaline, in small cloth sacks or pockets within each cabinet at the top. From time to time each few months during the year it is necessary to add more chemicals.

Mercuric chloride (bichloride of mercury) in solution is employed for poisoning botanical specimens. Before mounting, the specimen is dipped into this solution and then dried. Fumigation with carbon disulfide or a poisonous gas is another method. Or the specimens can be placed in a special cabinet or in an oven where insects may be killed with heat.

BOTANICAL TERMS

In order to use botanical references in the identification of trees it is necessary to understand the botanical terminology. The botanical descriptions for identification and also the natural classification are based principally upon morphology and anatomy. In distinguishing the species of trees differences in parts are used, such as presence or absence, number, arrangement, shape, size, etc. Thus, foresters should know the names of the important parts of a tree.

Flowering plants (Angiosperms), those with both flowers and seeds, and particularly the trees are composed of parts or organs. The principal organs are the root, stem, leaf, flower, and fruit. The first three, root, stem, and leaf, are vegetative in function and serve in the development and growth of the plant from the seed.

The root, composed usually of a vertical axis and branches, grows downward, anchors the plant in the soil, and absorbs the nutritive substances, but never bears leaves.

The stem, composed generally of a vertical axis and branches, grows upward (or opposite to the root), and supports the leaves and flowers.

The leaf, or foliar organ, generally is flat and green and functions to manufacture the food for the plant.

The other two organs, the flower and the fruit, are reproductive in function and serve in the production of seeds to propagate the species.

The flower, usually colored, is the organ of reproduction and consists of an axis or stem with modified leaves of four kinds: the calyx, corolla, the stamens with pollen, and the pistil with the feminine elements or the ovules.

The fruit is the mature ovary developed from the pistil and contains the mature seeds.

In the botanical classification of trees and also in the identification, the most important and most useful organs are the reproductive organs. The flower and fruit show the natural relations and botanical families much better than the other organs. The vegetative organs respond more easily and more rapidly to differences in the environment, such as climate and soil, and consequently are less constant and more variable. However, when a tree has neither flowers nor fruits, it is necessary to make use of these organs, such as leaves, bark, and wood, in the identification.

Systematic botanists have a special terminology of many words for describing the differences among the many thousands of species of plants. These words are derived from the ancient languages Latin and Greek. Many are similar to Spanish words and not difficult to remember. In the past centuries botanists wrote their books and descriptions in Latin, and for this reason there now remains for us a very rich botanical terminology. Actually there are more terms in plant taxonomy than are needed, and others are rarely used.

Foresters do not need as many terms as do botanists. In the study of dendrology we shall learn the common terms that are met with in descriptions of tropical trees. Other new or unfamiliar terms that are found in publications can be sought in botanical glossaries or other botanical references.

The root

The root generally is not important in classification of trees. As they are within the ground, roots are not seen nor easily examined. Also, they probably have very few characters useful in identification. However, a few species of trees have aerial roots which are useful in the forest for recognition.

The Bark

In the forest, the bark of tree trunks is used in identification by both the foresters and the country people. However, as the bark is not collected with botanical specimens, botanists usually are not familiar with the bark and do not use it in identification or classification.

Characters of the bark utilized in identification include external and internal color; the surface, texture, whether smooth, rough, or furrowed; if thin or thick; and if there is latex (colored juice or sap) or odor.

The trunk of some species has the base enlarged into buttresses, especially in the wet tropical forests.

It is difficult to describe the characters of bark and its differences. However, with experience in the forest one can recognize and distinguish various tree species by the bark.

The Stem

The stem bears the branches and leaves. The node is the point on the stem where one or more leaves are inserted. Sometimes it is enlarged or marked with a ring. The internode is the space on the stem extending between two consecutive nodes.

The bud is a short growing point of a stem, composed of young leaves or flowers and frequently also of some scales. In tropical regions generally there are no scales. However, in temperate regions with cold winters there are dormant buds covered with scales. There are two principal types of buds. The terminal bud is in the apex of the stem or

a branch. The lateral bud (or axillary bud) is located at the base of a leaf and can grow into a branch. (Branches, or lateral stems, are formed only at the nodes and from lateral buds.)

The leaf

A tree which has leaves in all seasons of the year is evergreen. If it remains without leaves during part of the year, such as in the dry or cold season, a tree is deciduous.

The parts of the leaf are the blade and the petiole. In some species there are also two (or one) stipules, or scales, usually deciduous, at the base. Also in some species the petiole is lacking, and the leaf is said to be sessile. Upon falling, the leaf makes a scar at the node.

Arrangement of leaves.--Alternate leaves are inserted separately or singly at different heights on the stem or one at a node. Opposite leaves are inserted by pairs at the same height on the stem or at a node. Verticillate leaves are inserted three or more at the same height on the stem or at a node.

In number of blades the leaves are simple and compound. The simple leaf consists of a single blade. The compound leaf is composed of usually several blades (two or more) which are called leaflets. Leaflets may have petioles or none but never have buds at the base. This character, the absence of the bud at the base, distinguishes a leaflet from a simple leaf. Compound leaves are divided into pinnate and digitate leaves. Leaves are pinnate when the leaflets are inserted along a common axis. If the axis is divided, the leaf can be twice pinnate (bipinnate) or three times pinnate (tripinnate). Leaflets are alternate in some species and opposite in others. If a pinnate leaf ends in a single leaflet it is odd pinnate (imparipinnate). If it has paired leaflets and ends in two leaflets it is even pinnate (paripinnate). Leaves are digitate (or palmate) when the leaflets are inserted together at the apex of the petiole.

Shapes of leaf blades.--The shape or form of leaf blades generally is characteristic of a species and is used in identification. There are many terms for describing the shape, but it is sufficient here to mention only the commonest types. A linear leaf has the blade long and narrow with margins parallel, for example in grasses or the grass family. A lanceolate leaf has the blade in the shape of a lance, several times longer than broad, broadest near the base, and pointed at the apex. (The reverse shape is called oblanceolate.) An ovate leaf has the blade oval but broadest toward the base, more or less as in the longitudinal section through an egg. (The reverse shape is called obovate.) An elliptic leaf has the blade oval but broadest at the middle. An oblong leaf has the blade several times longer than broad and with margins parallel. A circular leaf (or orbicular) has the blade more or less in a circle.

Margins of leaf blades.--The margin or border of the leaf blade can be: Entire, when the margin is smooth, or straight or curved but without teeth or lobes. Toothed, when it has pointed teeth, such as resembling a saw. Lobed, when it has deep indentations or lobes, which are larger than teeth but not deep enough to divide the blade into separate blades.

Apexes of leaf blades.--The apex of the leaf blade can be: Acuminate, gradually narrowed to the point in the shape of a wedge. Acute, with short point. Obtuse, or with blunt point. Rounded, or circular and without a point.

Bases of leaf blades.--The base of a leaf can be: Acute, with short point. Rounded, or circular and without a point. Heart-shaped, with a deep indentation as in the heart.

Venation of leaf blades.--According to the venation or network of veins, the leaf blade can be: Parallel-veined when the veins are parallel. Pinnate-veined or not-veined, when there is a single principal vein or midrib with other, lateral veins on the sides. Palmate-veined, when there are several principal veins which arise at the base and spread like fingers of the hand.

The Flower

A complete flower consists of four concentric parts, spirals or circles (whorls), called: the calyx, the corolla, the stamens (the androecium), and the pistil. However, some flowers lack one or more of these parts. The calyx is composed of modified small leaves generally green and called sepals. The corolla is composed of modified small leaves generally colored and called petals.

The stamens are the masculine organs of the flower and provide the pollen, which the male element. A stamen consists of two parts: the filament or stalk, generally thin, and the anther or enlarged part at the apex, composed of pollen sacs with the pollen grains.

The pistil is the feminine organ of the flower and is composed of one or more modified small leaves called carpels. The carpels form the ovary which is the feminine element and which is transformed into the fruit. The ovary contains one or more ovules which develop into the seeds. Other parts of the pistil are: the style, a small column above the ovary which supports the stigma, and the stigma, the uppermost enlarged part of the pistil, which receives the pollen grains. The stigma can have different shapes, such as rounded, lobed, or feathery.

The receptacle is the enlarged base of the flower where the floral parts are inserted. The peduncle is the stalk of the flower. If there are several flowers together, the stalk or secondary peduncle of each one is called a pedicel.

In the different families there are many differences in the shape of the flower and in the number, size, and arrangement of the parts. The parts of a circle or whorl can be separate (free) or united into a tube or united with other parts. The gamopetalous corolla has the petals united. The regular corolla has petals of equal size and symmetrically arranged. The irregular corolla has the petals unequal.

The carpel can be compared with a leaf on whose borders have developed the ovules. Also, the leaf has become incurved until the ovules are within a cavity of the ovary. The ovary encloses one or several small

cavities called cells or locules, which contain the ovules. An ovary formed from one carpel has necessarily a single cell. An ovary formed by two or more carpels has a single cell if the carpels are joined by their borders. If the carpels are folded toward the interior, they form as many cavities or cells as there are carpels. According to the number of cells the ovary can be: unilocular, bilocular, trilocular, or multilocular.

In position with respect to the other floral whorls, the ovary can be superior (free), or inferior (adhering). The superior ovary is free or separate in the middle of the flower and is inserted above the other whorls. The inferior ovary is united with other parts or whorls which appear to be inserted above.

Pollination is the transport of the pollen from the anther to the stigma. The pollen grain germinates and forms the pollen tube which carries the male element to the ovule. Fertilization is the union of the nucleus of the male element with the nucleus of the female element in the ovule to form the fertilized egg. The fertilized egg divides to form the embryo, and the ovule is transformed into the seed. The seed is the mature ovule and consists of the embryo with stored food and one or two seed coats.

The Inflorescence

The inflorescence is the arrangement or disposition of the flowers on the stem. It is terminal when it is at the apex of the stem or lateral (or axillary) when it is at the base of a leaf, on the side of the stem.

There are many types of inflorescences, but it is sufficient here to mention only some of the commonest. The solitary inflorescence (or simple) has the flowers isolated or one by one, separated by the leaves.

The spike has an elongated axis and flowers without pedicels. The raceme has an elongated axis and flowers with pedicels. The panicle is a compound raceme with the axis branched.

The umbel has the flowers together at the apex of the peduncle terminating spreading pedicels of equal length. (Also there is the compound umbel.) The head has the apex of the axis broadened as a disk and the flowers without pedicels. The cyme is a definite or determinate inflorescence with the principal axis ending in the first flower, and below arise other secondary axes with flowers.

The Fruit

The fruit develops from the mature ovary, contains the seeds, and sometimes bears other flower parts which persist. A simple fruit comes from a single pistil. An aggregate fruit develops from several pistils of a single flower (annona, strawberry, etc.). A multiple fruit comes from several flowers united (fig, pineapple, etc.).

Simple fruits are classified as dry and fleshy (juicy) and the dry fruits as indehiscent, if they do not open to discharge the seeds, or dehiscent, if they open.

There are several types of simple fruits. The akene is a dry indehiscent fruit with a single seed which does not adhere to the wall of the fruit. The nut is a dry indehiscent fruit with a single seed and with the wall of the fruit thick, hard, and woody. The pod (or legume) is a dry dehiscent fruit of one carpel which opens on two lines, while a follicle opens on one line. The capsule is a dry dehiscent fruit of two or more cells which opens on as many lines as there are cells.

The berry is a fleshy indehiscent fruit with several seeds. The drupe is a fleshy indehiscent fruit generally of one carpel (or more) and with a single seed (or more) enclosed in a stone or hard wall.

THE DISTRIBUTION OF TREES

Distribution of trees includes the geographic area where they occur, the climatic zones, the forest type, and the altitude or elevation above sea level.

In the study of plant geography there are two kinds of geographic distribution, that of the flora and that of the vegetation. The flora of a region is a list of the species of plants. The units of the flora are the species, genera, etc. The vegetation of a region is the appearance or the physiognomy of the plants or the growth form or habit of the commonest species. The units of vegetation are the plant communities, such as formations, associations, and forest types.

For example, let us suppose that two areas have only two species of plants, a tree species and a grass species. In one there is a single example of a tree and many individuals of the grass--it is a grassland. In the other there are many trees and few plants of the grass--it is a forest. Both areas have the same flora or list of species but have different vegetation. One is a grassland and the other a forest.

Thus, the distribution of the trees of a region can be studied from the viewpoint of the flora, or of the geographic range of each species. Distribution maps of each tree species can be made. Or from the viewpoint of vegetation vegetation types are studied in relation to climate. Maps of vegetation types such as forests, grasslands, etc., can be made. In dendrology both types of geographic distribution of plants are studied. Maps of the distribution of individual tree species and likewise of forest types are prepared.

Many tree species of tropical America are not sufficiently known for compiling maps of geographic distribution or analyzing the origin, the migrations, and the distribution. Some species have a broad range, while others are rare and local.

For example, some tree species have a very large distribution in tropical America from Central America or Mexico to Brazil. Those of ocean shores, such as the mangroves, have a wide range up to Florida in southeastern United States. Other trees are confined to high mountains, still others to grassy plains or to deserts, etc. Few are local or endemic. A job for the future is to prepare more detailed distribution maps.

The study of vegetation and distribution of vegetation types is often included in a subdivision of biology known as ecology. Much of the following material treated under ecology might equally well be considered as forestry or dendrology.

ECOLOGY

The name ecology (originally spelled oecology) was first used in 1869 by the German zoologist E. Haeckel. It is derived from two Greek roots meaning home and discourse or study, or literally, the study of homes. Ecology is defined as the study of organisms in relation to their environment. Opinions differ as to whether ecology really is a subject matter branch of biology or whether it is merely a point of view of studying biology. Thus, ecology might be biology with emphasis on the surroundings of animals and plants.

Ecology is often subdivided into autecology and synecology. The former prefix is a Greek root meaning self, while the latter means together. Autecology is the ecology of the individual, or study of an individual organism or species in relation to its environment. By some authors autecology is considered as physiology. For example, an investigation of a single tree species in relation to its surroundings. In contrast, synecology is the ecology of the group, or study of groups of organisms, or communities in relation to their environment.

As the subject matter or viewpoint of ecology is so large that it embraces biology, ecology is usually divided into animal ecology, which treats of animals in relation to their environment, and plant ecology, which deals with plants in relation to their environment. By some authors plant ecology is restricted to synecology as the study of plant communities or vegetation.

Plant ecology was established by the Danish botanist E. Warming in his classic book, Oecology of Plants, first published in Danish in 1895. Another important early reference was Plant Geography upon a physiological Basis, by the German botanist A. F. W. Schimper, which appeared in German in 1898.

The environment, or surroundings, includes everything that may affect an organism or group of organisms in any way. In plant ecology one classification divides the environment into factors, or environmental factors, as follows: (1) climatic factors, or climate, which includes temperature, rainfall, and other widely distributed elements chiefly from the atmosphere; (2) physiographic or edaphic factors, such as topography and soil; (3) biotic factors, or other organisms. Because modern man has had such a profound influence, mostly destructive, on other organisms, man has often been separated from the third group as a fourth group of factors called anthropeic factors. As fire has become widespread and has serious effects upon vegetation, it is often listed as a fifth group, known as pyric factors.

One of the principles of ecology is often called balance of life, or web of life. This means that organisms living together in a group or community are more or less balanced or at equilibrium, interrelated by

their mutual requirements for food for energy, the original source of which is the sun. The activities within the group and particularly the number or individuals are in balance with the available supply of energy. A simple relationship of several kinds of organisms, each dependent upon a surplus of another for food, is called a food chain. For example, in a pond the green algae or pond scums make food with energy received from the sun and are eaten by microscopic animals. These in turn are consumed by less minute aquatic animals later eaten by small fish. The latter are devoured by larger fish, which are prey of birds, mammals, and man. The relationship of foods generally is more complex with branches and is a network or web rather than a chain.

Adaptation, or the adjustment of organisms to their environment, is another principle of ecology. Each species fits a set of external conditions, sometimes within very narrow limits, where it is in equilibrium or in balance.

An important principle of ecology is that of change, also known as the dynamic viewpoint. The whole universe, or rather the universes about us, from the giant galaxies of stars to the minute particles within the atom, is changing or in motion all the time. Specifically, the climate of a region is slowly changing and the soil is changing. Therefore, the vegetation, which is dependent upon climatic and soil factors, is slowly changing at a particular place too.

Classification based on water relations.--Tree species as well as other plants and also vegetation can be classified on the basis of water relations or distribution based upon rainfall or water. There are three or more principal groups. A hydrophyte is a plant of wet places with adequate water, such as one growing along river banks. A xerophyte is a plant of dry places with very little water as in deserts. A mesophyte is a plant of places with average amounts of water, such as forests neither very humid nor very dry. Most trees and most other plants belong here.

A tropophyte, literally change plant, is a plant which grows under conditions which change markedly during the year, such as plants of places where the quantity of water changes during the year from much to little in seasons from humid to dry. For example, a deciduous tree in a tropical savanna. The term tropophyte also is applied in the temperate zones where the temperature changes during the year to a tree which is deciduous in the cold or winter season.

A halophyte is a plant of places with concentrations of salt, such as sea shores and alkali deserts. For example, the mangroves.

Also for vegetation there are used the adjective terms hydrophilous, xerophilous, mesophilous, tropophilous, and halophilous. However, this Greek suffix meaning loving and suggests teleology, and old idea that there is a cause or reason for everything. Trees do not have intelligence and cannot love water. In biology generally and particularly in ecology one must be careful not to seek a purpose underlying each adaptation or peculiar structure.

Plant communities.--In the vegetation the plants arrange themselves in groups or communities. Any group of plants in a place or common habitat can be called a plant community. This term has no restrictions in size.

A plant formation is a plant community of the highest rank and characterized by a definite appearance or physiognomy and composed of plants with a definite growth form. For example, the rain forest or the páramo.

The formation is composed of smaller units, the associations. A plant association is a division of a formation with a definite floristic composition and with certain dominant species. The forest types of the foresters correspond to the plant associations of the ecologists. For example, a forest of Podocarpus. Some associations are limited in extent by climatic factors and others, the edaphic associations, by the factors of the soil.

Plant communities are constantly changing though sometimes very slowly. Great changes in the composition of a community follow changes in the environment which in turn may be caused by the organisms living there. The term succession, or plant succession, is given to the changes in the plant communities, one following another, at the same place. The succession leads to the climax association, which is the highest, most complex, and most nearly stable community existing in the climate of that area. Where the highest vegetation of the succession is limited by edaphic factors rather than climate, the climax association, such as on a particular soil type, is called an edaphic climax. The mangrove swamp forest is an example.

Plant successions are of two main kinds, primary and secondary. A primary succession begins on a bare area previously without vegetation, such as a new island, a new volcano, or landslide. A xerarch succession, or xerosere, is one beginning with a dry bare habitat and leading to a climax. A hydrarch succession, or hydrosere, starts with a wet habitat and continues to a climax. The climax has medium moisture conditions, rather than extreme. Virgin forests represent climaxes or normal developmental stages.

A secondary succession is that following a disturbance in the normal, primary succession. For example, after cultivation, cutting of forests, fire, overgrazing, or wind damage. In densely settled regions most of the vegetation has been disturbed by man's activities. Few areas of virgin forests remain and the forests are largely secondary. Ultimately the secondary succession leads to the same climax as the primary succession, but the rate depends upon the degree of disturbance. If great, such as when soil formed during thousands of years is eroded away, the recovery is very slow.

Groups of plant formations of the world.--The plant formations of the world can be arranged into four or five large groups:


1. Forests, characterized by trees.
2. Grasslands, characterized by grasses.
3. Deserts, characterized by shrubs, spiny trees, cacti, etc., and by scarcity of water.
4. Tundras, characterized by lichens, mosses, etc. and by very low temperatures.
5. Oceans, characterized by seaweeds or algae.

LEAF CHARACTERISTICS FOR TREE IDENTIFICATION

(CARACTERISTICAS DE LAS HOJAS PARA LA IDENTIFICACION DE LOS ARBOLES)


Leaf Shapes

(Forma de las Hojas)



1	Acicular	Acicular
2	Subulate	Alesnada
3	Linear	Linear
4	Oblong	Oblonga
5	Lanceolate	Lanceolada
6	Oblanceolate	Oblanceolada
7	Ovate	Aovada
8	Obovate	Obovada
9	Elliptical	Elíptica
10	Oval	Ovalada
11	Orbicular	Orbicular
12	Reniform	Reniforme
13	Cordate	Cordiforme
14	Deltoid	Deltoide
15	Rhomboid	Romboide
16	Spatulate	Espatulada


Leaf Margins
(Borde de las Hojas)



- | | | |
|----|----------------|---------------------|
| 1 | Revolute | Enrollada |
| 2 | Entire | Entera |
| 3 | Repand | Ondulada |
| 4 | Sinuate | Sinuada |
| 5 | Crenate | Crenada |
| 6 | Serrate | Aserrada |
| 7 | Dentate | Dentada |
| 8 | Doubly serrate | Doblemente aserrada |
| 9 | Doubly crenate | Doblemente crenada |
| 10 | Lobed | Lobulada |
| 11 | Cleft | Hendida |
| 12 | Parted | Partida |

Apices


(Apices)



- | | | |
|---|------------|------------|
| 1 | Acuminate | Acuminada |
| 2 | Acute | Aguda |
| 3 | Mucronate | Mucronada |
| 4 | Cuspidate | Cuspidada |
| 5 | Obtuse | Obtusa |
| 6 | Rounded | Redondeada |
| 7 | Truncate | Truncada |
| 8 | Emarginate | Emarginada |

Bases

(Bases)



- | | | |
|----|---------------|------------|
| 9 | Cuneate | Cuneada |
| 10 | Acute | Aguda |
| 11 | Cordate | Cordada |
| 12 | Inequilateral | Obliqua |
| 13 | Obtuse | Obtusa |
| 14 | Rounded | Redondeada |
| 15 | Truncate | Truncada |
| 16 | Auriculate | Auriculada |

